

Direct Oral Anticoagulants (DOACs) Guide #2

Brand Name	Generic Name	Strength	GoodRx Price/ Unit	GoodRx Price/ 30 Days
Anticoagulant				
Coumadin	warfarin (once daily)	1mg	\$ 0.13	\$4.00/30
		2mg	\$ 0.13	\$4.00/30
		2.5mg	\$ 0.13	\$4.00/30
		3mg	\$ 0.13	\$4.00/30
		4mg	\$ 0.13	\$4.00/30
		5mg	\$ 0.13	\$4.00/30
		6mg	\$ 0.13	\$4.00/30
		7.5mg	\$ 0.13	\$4.00/30
		10mg	\$ 0.13	\$4.00/30
Warfarin – Reversal Agent				
Mephyton	phytonadione (vitamin K)	5mg	-	\$20.66/1 dose
Kcentra	prothrombin complex concentrate	Individualize dosing based on current pre-dose INR Administer with vitamin K	-	\$1,685.41 (500-unit vial x 1)
Factor Xa Inhibitors				
Eliquis	apixaban (twice daily)	2.5mg	\$ 8.76	\$525.47 /60
		5mg	\$ 8.76	\$525.47/60
Xarelto	rivaroxaban (once or twice daily)	10mg	\$ 17.11	\$513.37/30
		15mg	\$ 17.11	\$513.36/30
		20mg	\$ 17.44	\$523.11/30
Savaysa	edoxaban (once daily)	15mg	\$ 12.95	\$388.64/30
		30mg	\$ 13.19	\$395.71/30
		60mg	\$ 13.64	\$409.30/30
Thrombin Inhibitor				
Pradaxa	dabigatran (once or twice daily)	75mg	\$ 8.09	\$485.43/60
		110mg	\$ 8.07	\$484.43/60
		150mg	\$ 8.11	\$486.43/60

Factor Xa – Reversal Agent						
Andexxa	andexanet alfa	Usual dose 400mg or 800mg – dosing is dependent on FXa inhibitor used, dose and timing since last patient dose	-	-	-	\$26,400.00 (4 x 200mg vials)
Thrombin Inhibitor – Reversal Agent						
Praxbind	idarucizumab injection	5g/100mL	-	-	-	\$5,340.52 (2 x 2.5mg vials)

Items in green are available as generics. Medications in red are available as brands only

Sources: AmerisourceBergen ABC Passport and GoodRx (GRx) – accessed 9/2/2022

Perioperative/Periprocedural Management

Pradaxa (dabigatran)	Eliquis (apixaban)	Xarelto (rivaroxaban)	Savaysa (edoxaban)	Coumadin (warfarin)
<p>CrCl ≥50 mL/minute: Hold 1 to 2 days before surgery</p> <p>CrCl <50 mL/minute: Hold 3 to 5 days before surgery</p> <p>May consider holding for >5 days in patients undergoing major surgery, spinal puncture, or insertion of a spinal or epidural catheter or port</p>	<p>Hold at least 24 to 48 hours, depending on risk or location of bleeding, before elective surgery or invasive procedures</p>	<p>Hold at least 24 hours before surgery; longer duration of treatment cessation may be necessary based on individual patient situation and physician clinical judgment</p>	<p>Discontinue at least 24 hours prior to elective surgery or invasive procedures</p>	<p>Hold at least 5 days before surgery; depending on urgency of surgery/procedure, may administer low-dose IV or oral vitamin K</p> <p>Minor dental and minor dermatological procedures or cataract surgery: Continue warfarin (with hemostatic agent [dental] or local hemostasis [dermatological]); may also discontinue use 2 to 3 days prior to dental procedures</p> <p>Patients with prior stroke undergoing dental procedures should routinely continue warfarin</p>

Resource: Lexicomp – Accessed 8/22/19